arXiv:1210.6073v1 [cond-mat.stat-mech] 22 Oct 2012

Stokes–Einstein relation violation and the upper critical dimension of the glass transition

Patrick Charbonneau,[1,2] Giorgio Parisi,[3] and Francesco Zamponi[4]

[1]*Department of Chemistry, Duke University, Durham, North Carolina 27708, USA*
[2]*Department of Physics, Duke University, Durham, North Carolina 27708, USA*
[3]*Dipartimento di Fisica, Sapienza Universitá di Roma, INFN,*
Sezione di Roma I, IPFC – CNR, P.le A. Moro 2, I-00185 Roma, Italy
[4]*LPT, Ecole Normale Supérieure, CNRS UMR 8549, 24 Rue Lhomond, 75005 France*

Using numerical simulations we investigate the dimension evolution of the Stokes–Einstein relation violation in a simple glass forming system. The results strongly support the existence of an upper critical dimension $d_u = 8$ above which the Stokes-Einstein relation is not violated, as predicted by the mean-field theory of glasses. In addition, the associated critical exponents evolve linearly in $8-d$ below $d = 8$, which suggests that a suitable renormalization group expansion around the mean-field theory could provide a quantitative description of the glass problem. An analysis of single-particle motions in terms of Lévy flights offers a possible scenario for relating these corrections to the mean-field description with the dynamical heterogeneity of glass formers.

The glass problem is notoriously difficult. Constructing a completely satisfying theory for how rapidly a fluid turns sluggish with only unremarkable structural changes remains hotly debated [1]. A key hurdle is that the theories of the glass transition typically provide insufficiently precise predictions for decisive experimental or numerical tests to be performed. Identifying the "smoking gun" that could tell theories apart has thus long been a priority.

A promising way of addressing this issue is to investigate the glass transition as a function of spatial dimension d, which was recently achieved in numerical simulations [2]. In low d various phenomena kinetically compete in glass-forming fluids: *(i)* crystal nucleation, *(ii)* barrier hopping due to thermal activation, and *(iii)* trapping in phase space due to the proximity of ergodicity breaking. Different theories give more or less weight to these physical processes, which are so well enmeshed that they are typically hard to tell apart. An advantage of increasing d is that both nucleation and barrier hopping get strongly suppressed. Attention can then be focused on the onset of ergodicity breaking [2].

A central quantity on which most theories give specific predictions is the violation of the Stokes–Einstein relation (SER). According to the SER the structural relaxation time τ_α, which is proportional to the viscosity in a dense fluid, and diffusivity D should give a constant $D\tau_\alpha$. Yet in d=2 and 3, a large violation of this relation is observed upon approaching the glass transition [3–6]. In most cases it is found that $D \propto \tau_\alpha^{-1+\omega}$ [7–9]. The SER violation is traditionally attributed to a large spatial heterogeneity of the "local relaxation times" [9]. Diffusion is dominated by the faster regions, while viscosity and τ_α are dominated by the slower regions. Hence, $1/D \ll \tau_\alpha$ and one expects $\omega \geq 0$ [9], as is indeed observed numerically [6–8].

Explanations for the evolution of the SER violation with d fall into two categories. First, theories based on adapting the standard description of critical phenomena to the glass problem predict an upper critical dimension d_u. For $d > d_u$ a mean-field theory is expected to give the correct critical description, while for $d < d_u$ critical fluctuations qualitatively change the behavior of the system by renormalizing the scaling exponents or eliminating the transition altogether. In this scenario, SER violations would disappear for $d \geq d_u$, and it is expected that $d_u = 8$ based both on static [10, 11] and dynamical [9] descriptions. In particular, a scaling argument proposed in Ref. [9] gives $\omega \sim (8-d)/(4\gamma)$, where γ is the exponent that describes the divergence of the relaxation time with density (and, more generally, temperature) near the onset of ergodicity breaking [2]. Second,



FIG. 1: *Main panel*: The self-diffusion D and the structural relaxation time τ_α are related by a power-law with exponent $-1+\omega$ (dashed lines), where ω changes with d. Note that the power-law regime is truncated at low d by activated events on a scale similar to that observed for other dynamical quantities [2]. Over the limited remaining dynamical rang for d=3, ω=0.19(3). *Inset*: The exponent ω (symbol) linearly vanishes as $\omega = A(d_u - d)$ with $d_u = 8$ and $A = 0.038(3)$ (solid line).



FIG. 2: *Main panel*: As the system becomes more sluggish, $P(r,\tau_\alpha)$ (here for d=4) develops a long tail that is partly hidden by structural oscillations on a scale $r \sim 1$. *Inset*: As dimension increases, the tail of $P(r,\tau_\alpha)$ for isodiffusive systems ($2dD = 0.0008(3)$) gradually disappears.

other theories, e.g., those based on a certain class of kinetic models [12], predict that the critical behavior of the system is qualitatively similar in all spatial dimensions, hence $d_u = \infty$ [7]. In the East model studied in Refs. [7, 12, 13], for instance, it is found that $\omega \approx 0.2$ for all d. The existence of d_u can thus be directly established by studying the SER violation as a function of d. An attempt in this direction was done in Refs. [8, 14]. In this letter, we perform numerical simulations to measure ω in higher dimensions, and find that $d_u = 8$. We also rationalize that the SER violation can be interpreted in terms of a Lévy flight that describes the underlying behavior of the particle displacement distribution on the scale of τ_α.

Simulations – Event-driven molecular dynamics simulations of hard spheres in dimension d=3–9 are performed, following the method of Refs. [2, 15, 16] for at least $N = 8000$ identical particles of unit diameter σ, except for d=3 where an equimolar binary mixture with a diameter ratio 7:5 is used to prevent crystallization [17, 18]. Simulations are run at constant unit inverse temperature β with time t expressed in units of $\sqrt{\beta m \sigma^2}$ for particles of unit mass m. The diffusivity is extracted from the long-time behavior of the mean-square displacement $D = \lim_{t\to\infty}\langle r^2\rangle_t/2dt$, where $\mathbf{r} = \mathbf{r}(t) - \mathbf{r}(0)$, and τ_α is the characteristic decay time of the self-intermediate scattering function $F_s(k^*,t) = \langle e^{i\mathbf{k}^*\cdot\mathbf{r}}\rangle$ at the wave vector k^* that maximizes the first peak of the structure factor. Figure 1 confirms that upon approaching the glass transition the regime where $D \propto \tau_\alpha^{-1+\omega}$ extends with dimension, in agreement with activated relaxation processes getting increasingly suppressed. The results are also consistent with ω vanishing linearly at $d_u = 8$, which indicates that a finite upper critical dimension indeed exists. However, we find $\omega \sim 0.038(8-d)$, while the analysis of Ref. [9] gives $\omega \sim 0.12(8-d)$, using earlier evaluations of $\gamma \approx 2.1$ [2]. This discrepancy calls for a better investigation of the physical origin of the SER violation for $d < d_u$.

Both D and τ_α are themselves functionals of the self van Hove function $G_s(r,t)$ [19]. Because D weighs large displacements more than τ_α, the SER violation thus suggests that $G_s(r,\tau_\alpha)$ has an anomalous behavior at large r. The spherically integrated displacement distribution $P(r,\tau_\alpha) = \Omega_d r^{d-1} G_s(r,\tau_\alpha)$ indeed develops a fat tail at large r when τ_α increases (Fig. 2), as was previously reported in various d=2 and 3 systems [18, 20, 21]. This behavior is remarkably distinct from the Gaussian distribution expected in the long-time diffusive regime [19], and is also different from the exponential regime that is observed in the caging, or β, regime [2]. In the structural relaxation regime, the fat tail of $G_s(r,\tau_\alpha)$ results in a larger than expected mean square displacement and in the SER violation. By contrast, with increasing d the bulging tail resorbs and the SER is gradually recovered (Fig. 2). To further substantiate this claim, we introduce a time scale τ_D, such that $\langle r^2\rangle_{\tau_D} = 2d/(k^*)^2$. Numerically, we find that *(i)* τ_D corresponds to the beginning of the diffusive regime and thus $\langle r^2\rangle_t \sim 2dDt$ for $t > \tau_D$, and *(ii)* $D \propto \tau_D^{-1}$ (Fig. 3). The SER violation and *(ii)* imply that $\tau_D \sim \tau_\alpha^{1-\omega} \ll \tau_\alpha$. Hence from *(i)* we deduce $\langle r^2\rangle_{\tau_\alpha} \sim 2dD\tau_\alpha \propto \tau_\alpha/\tau_D \propto \tau_\alpha^\omega$, i.e. the second moment of $G_s(r,\tau_\alpha)$ diverges with τ_α. This argument establishes the relation between the SER violation and the fat tail of the self van Hove function, as illustrated in Fig. 3.

Lévy Flight Model – A minimal model for describing the physical consequences of this distribution anomaly can be formulated in terms of a truncated Lévy flight [22–24], which amount to the following two assumptions. *(a)* For large τ_α, the distribution $P(r,\tau_\alpha)$ converges to a finite limit around its maximum $r_{\rm typ}(\tau_\alpha) \sim 1/k^*$ that is almost independent of τ_α (Fig. 2), while its behavior for $r \gg r_{\rm typ}(\tau_\alpha)$ is described by

$$P(r,\tau_\alpha) = \frac{r_{\rm typ}(\tau_\alpha)^\mu}{r^{1+\mu}}\mathcal{P}[r/\xi(\tau_\alpha)] \ , \qquad (1)$$

where we defined a dynamical length $\xi \sim \tau_\alpha^{1/\zeta}$ and the critical exponents μ and $\zeta > 0$. In this model $\mathcal{P}(0)$ is a constant and $\mathcal{P}(x)$ rapidly vanishes at large x. We assume that $\mu < 2$ for $d < 8$, in order for the second moment of $P(r,\tau_\alpha)$ to diverge when $\tau_\alpha \to \infty$. Note that the bounded limit distribution at small r guarantees its normalizability, which is consistent with the majority of displacements taking place on a scale $r_{\rm typ}(\tau_\alpha)$ at τ_α. *(b)* On a time scale $t \gg \tau_\alpha$, particle displacements are described by a random walk with uncorrelated elementary steps distributed according to $P(r,\tau_\alpha)$. Stated differently, the self van Hove function at $\tau = t/\tau_\alpha > 1$ can be obtained by convolutions of the function evaluated at $\tau = 1$, and thus

$$F_s(k,t) \sim e^{-\tau f_{\tau_\alpha}(k^2)} \ . \qquad (2)$$



FIG. 3: *Upper panel:* The mean square displacement $\langle r^2\rangle_t = R^2(2,t)$ as a function of time for $d=5$ and several densities upon approaching the glass transition. Crosses indicate the time τ_D such that $\langle r^2\rangle_{\tau_D} = 2d/(k^*)^2$ (long-dashed line), which marks the beginning of the diffusive regime. Asterisks indicate the relaxation time τ_α, which shows the growth of $\langle r^2\rangle_{\tau_\alpha} \propto \tau_\alpha^\omega$ (short-dashed line). *Inset:* The separate evolution of τ_α (asterisks) and τ_D (crosses) with D, where $D \sim \tau_D^{-1}$ (long-dashed line) and $D \sim \tau_\alpha^{-1+\omega}$ (short-dashed line). *Lower panel:* $R^2(z,t)$ for z=0.1, 2, and 4 in d=5 for $\varphi = 0.268$ (squares). The Lévy flight regime extends from $t > \tau_\alpha$ to the rightmost part of the figure, where the functions cross over to the Gaussian diffusive regime. The predictions of Eq. (7) are consistent with the simulation results. The dot-dashed line has slope $2/\mu$ with $\mu = 1.423$, following Eq. (6), while the full line has slope 1. The non-Gaussian parameter $\alpha_2(t)$ (circles) shows that non-Gaussianity is maximal at τ_D and persists across the Lévy flight regime, while $\alpha_2(t) \sim 1/t$ at larger times in the Gaussian diffusive regime.



FIG. 4: *Main panel*: The value of $\alpha_2(\tau_\alpha)$ versus τ_α for several dimensions indicates that $\alpha_2(\tau_\alpha) \sim \tau_\alpha^\eta$ over the non-activated regime. *Inset*: The exponent η (symbol) scales linearly $\eta = A(d_u - d) + \eta_\infty$ with $d_u = 8$, $A = 0.017(2)$, and $\eta_\infty = 0.23(1)$ (solid line), which suggests that $\eta \approx 0.315$ in d=3.

If these two assumptions are true, at large times particles perform a random walk whose distribution of elementary steps acquires a power-law tail at the onset of ergodicity breaking, i.e. when the structural relaxation time diverges. This Lévy flight description of the displacement tail at τ_α offers a natural microscopic explanation for SER violation. Numerically, however, the large r tail of $P(r,\tau_\alpha)$ is modulated by structural oscillations that partially obscure the presumed power-law behavior (see Fig. 2). More robust consequences of the model must thus be examined to assess its relevance.

Exponents and Scaling – Considering the fractional moments of the displacements, defined as $R^2(z,t) = [\langle r^z\rangle_t]^{\frac{2}{z}}$, one can show from Eq. (1) that

$$R^2(z,\tau_\alpha) \sim \begin{cases} r_{\rm typ}(\tau_\alpha)^{2\mu/z}\xi(\tau_\alpha)^{2(1-\mu/z)} & \text{for } z > \mu \ , \\ r_{\rm typ}(\tau_\alpha)^2 C(z) & \text{for } z < \mu \ , \end{cases} \tag{3}$$

where $C(z)$ remains finite for $\tau_\alpha \to \infty$ (but diverges for $z \to \mu$). Using this form, and recalling that $\mu < 2$, we obtain that

$$\langle r^2\rangle_{\tau_\alpha} = R^2(2,\tau_\alpha) \sim \xi(\tau_\alpha)^{2(1-\mu/2)} \sim \tau_\alpha^{2(1-\mu/2)/\zeta} \ . \tag{4}$$

Hence, $\omega = 2(1-\mu/2)/\zeta$ and the non-Gaussian parameter scales as

$$\alpha_2(\tau_\alpha) = \frac{d}{d+2}\frac{\langle r^4\rangle_{\tau_\alpha}}{\langle r^2\rangle_{\tau_\alpha}^2} - 1 \sim \frac{[R^2(4,\tau_\alpha)]^2}{[R^2(2,\tau_\alpha)]^2} \sim \tau_\alpha^{\mu/\zeta} = \tau_\alpha^\eta \ . \tag{5}$$

Numerical results give $\eta \sim 0.23(1) + 0.017(2)(8-d)$ (Fig. 4), so from the measured $d \leq 8$ exponents we deduce that

$$\begin{aligned} \mu &= \frac{2\eta}{\eta+\omega} \approx \frac{0.46 + 0.034(8-d)}{0.23+0.055(8-d)} \ , \\ \frac{1}{\zeta} &= \frac{\eta+\omega}{2} \approx \frac{1}{2}[0.23 + 0.055(8-d)] \ , \end{aligned} \tag{6}$$

which gives $\mu \approx$1.25 and $1/\zeta \approx$0.25 in d=3.

From Eq. (2) we also obtain that for finite τ_α there exists a “Lévy flight” regime $1 \leq \tau \ll \xi^\mu$ in which

$$R^2(z,t) \sim \begin{cases} r_{\rm typ}(\tau)^{2\mu/z}\xi^{2(1-\mu/z)} & \text{for } z > \mu \ , \\ r_{\rm typ}(\tau)^2 C(z) & \text{for } z < \mu \ , \end{cases} \tag{7}$$

with a typical displacement $r_{\rm typ}(\tau) \sim \tau^{1/\mu}$. When $\tau \sim \xi^\mu$, and thus $t \sim \tau_\alpha \xi^\mu \sim \tau_\alpha^{1+\eta}$, we obtain $R^2(z,t) \sim r_{\rm typ}(\xi^\mu)^2 \sim \xi^2$ for all z. From that point on, i.e. for $t > \tau_\alpha^{1+\eta}$, the two length scales $r_{\rm typ}(\tau)$ and ξ merge and one recovers the trivial Gaussian diffusive regime. The curves for all z then merge and scale as τ. Note that Eq. (1) is valid for $\tau_\alpha < t < \tau_\alpha^{1+\eta}$, with a t-dependent $r_{\rm typ}$. Although our results are well captured by keeping ξ constant, we cannot exclude the possibility of a small time dependence of ξ in this regime.

Figure 3 shows that $R^2(z,t)$ follows the prediction of Eq. (7) over one decade of t in $d = 5$, which strongly suggests that the postulated model is accurate. Because we can only access a limited dynamical range and the exponent η is quite small, the Lévy flight regime $\tau_\alpha < t < \tau_\alpha^{1+\eta}$ is, however, rather limited and therefore a complete verification of Eq. (7) is not yet possible. The key difficulty is that the range of accessible τ_α is always limited. Activated events interfere with the slowdown in d=3–5, and numerical simulations of denser systems in $d \geq 6$ are currently beyond computational reach.

Conclusion – In summary, the critical behavior of the SER violation disappears for $d \geq 8$, which indicates that the glass transition has an upper critical dimension $d_u = 8$. For $d \geq d_u$ the mean-field theory of the glass transition is thus a good description, and extending this theory using renormalization group methods should top the research agenda, as suggested in Ref. [9]. In particular, a better understanding of the physical behavior below d_u would be helpful to clarify why the prediction of Ref. [9] for the exponent ω is not quantitatively accurate.

Of particular interest in this respect is understanding how dynamical heterogeneity, which is well documented in the literature [8, 18, 25] and is confirmed by a preliminary analysis of our data in all dimensions, results in a fat displacement tail at τ_α. Our analysis of particle displacements only covers the regime $t \geq \tau_\alpha$, so we still lack a complete understanding of the microscopic mechanism on scales $t < \tau_\alpha$. The question is especially interesting around τ_D, where diffusion sets in and non-Gaussianity is maximal. A possible explanation for the Lévy flight-type distributions is for displacements on shorter timescales to be random in time, but correlated in space, resulting in large spatial jumps [24]. Because this process is necessarily collective, the description also offers a direct link between the SER violation and a growing dynamical heterogeneity [24].

A related point that deserves further examination is the connection between the dynamical length ξ from the Lévy distribution expression in Eq. (1) and the spatial extension of dynamical heterogeneity ξ_4, as quantified by four-point correlations [8, 18, 25]. An argument proposed in Ref. [24] relates these two lengths in a different dynamical regime, but it might be possible to extend the argument to the regime considered here. Mean-field treatments indeed predict that $\xi_4 \sim \tau_\alpha^{1/(4\gamma)}$ for $d \geq 8$ [10, 11, 26], which, using $\gamma = 2.1$ [2], gives $\xi_4 \sim \tau_\alpha^{0.12}$. This scaling is remarkably close to the one we obtain here for the dynamical length, i.e. $\xi \sim \tau_\alpha^{1/\zeta} = \tau_\alpha^{0.115}$ in $d = 8$.

Acknowledgements – We thank J.-P. Bouchaud and D. R. Reichman for very useful discussions. PC acknowledges NSF support No. NSF DMR-1055586. GP acknowledges financial support from the European Research Council through ERC grant agreement No. 247328.

[1] G. Tarjus, in *Dynamical Heterogeneities and Glasses*, edited by L. Berthier, G. Biroli, J.-P. Bouchaud, L. Cipelletti, and W. van Saarloos (Oxford University Press, 2011).
[2] P. Charbonneau, A. Ikeda, G. Parisi, and F. Zamponi, Proc. Nat. Acad. Sci. U. S. A. **109**, 13939 (2012).
[3] F. H. Stillinger and J. A. Hodgdon, Phys. Rev. E **50**, 2064 (1994).
[4] G. Tarjus and D. Kivelson, J. Chem. Phys. **103**, 3071 (1995).
[5] I. Chang and H. Sillescu, J. Phys. Chem. B **101**, 8794 (1997).
[6] D. N. Perera and P. Harrowell, Phys. Rev. Lett. **81**, 120 (1998).
[7] L. Berthier, D. Chandler, and J. P. Garrahan, Europhys. Lett. **69**, 320 (2005).
[8] J. D. Eaves and D. R. Reichman, Proc. Nat. Acad. Sci. U.S.A. **106**, 15111 (2009).
[9] G. Biroli and J.-P. Bouchaud, J. Phys.: Cond. Matt. **19**, 205101 (2007).
[10] S. Franz, G. Parisi, F. Ricci-Tersenghi, and T. Rizzo, Eur. Phys. J. E **34**, 1 (2011).
[11] S. Franz, H. Jacquin, G. Parisi, P. Urbani, and F. Zamponi, `arXiv:1206.2482` (2012).
[12] A. S. Keys, L. O. Hedges, J. P. Garrahan, S. C. Glotzer, and D. Chandler, Phys. Rev. X **1**, 021013 (2011).
[13] D. Ashton, L. Hedges, and J. P. Garrahan, J. Stat. Mech. **2005**, P12010 (2005).
[14] P. Charbonneau, A. Ikeda, J. A. van Meel, and K. Miyazaki, Phys. Rev. E **81**, 040501 (2010).
[15] M. Skoge, A. Donev, F. H. Stillinger, and S. Torquato, Phys. Rev. E **74**, 041127 (2006).
[16] P. Charbonneau, A. Ikeda, G. Parisi, and F. Zamponi, Phys. Rev. Lett. **107**, 185702 (2011).
[17] L. Berthier and T. A. Witten, Phys. Rev. E **80**, 021502 (2009).
[18] E. Flenner, M. Zhang, and G. Szamel, Phys. Rev. E **83**, 051501 (2011).
[19] J.-P. Hansen and I. R. McDonald, *Theory of simple liquids* (Academic Press, London, 1986).
[20] P. Chaudhuri, L. Berthier, and W. Kob, Phys. Rev. Lett. **99**, 060604 (2007).
[21] C. Heussinger, L. Berthier, and J.-L. Barrat, Europhys. Lett. **90**, 20005 (2010).
[22] J.-P. Bouchaud and A. Georges, Phys. Rep. **195**, 127 (1990).
[23] I. Koponen, Phys. Rev. E **52**, 1197 (1995).
[24] F. Lechenault, R. Candelier, O. Dauchot, J.-P. Bouchaud, and G. Biroli, Soft Matter **6**, 3059 (2010).

[25] L. Berthier, G. Biroli, J.-P. Bouchaud, L. Cipelletti, and W. van Saarloos, eds., *Dynamical Heterogeneities and Glasses* (Oxford University Press, 2011).

[26] G. Biroli, J.-P. Bouchaud, K. Miyazaki, and D. R. Reichman, Phys. Rev. Lett. **97**, 195701 (2006).